



HAVE YOU SCHEDULED YOUR CHILD'S NEXT WELL-CHILD CARE APPOINTMENT?



WELL-CHILD VISITS

- 3-5 days (for breast-fed newborns)
- 1-2 weeks
- 1 month
- 2 months
- 4 months
- 6 months
- 9 months
- 12 months
- 15 months
- 18 months
- 24 months (2 years old)
- Annually for ages 3-21



SCREENINGS

- **Hearing:** 4-6 years
- **Vision:** 3-6 years



IMMUNIZATIONS

- **Birth:** Hepatitis B #1
- **2 weeks:** No immunizations
- **1 month:** No immunizations
- **2 months:** Vaxelis #1: PCV-13 #1, Rotarix #1
- **4 months:** Vaxelis #2: PCV-13 #2, Rotarix #2
- **6 months:** Vaxelis #3: PCV-13 #3, Influenza #1
- **9 months:** Influenza #2: Lead, Hemoglobin + Hematocrit
- **12 months:** MMR #1, Varicella #1, Hepatitis A #1
- **15 months:** DTaP #4, Hib #4, and PCV-13 #4
- **18 months:** Hepatitis A #2 (6 months after 1st dose)
- **2 years:** No immunizations. Lead, Hemoglobin + Hematocrit
- **3 years:** No immunizations. Lead, Hemoglobin + Hematocrit, TB Screening
- **4 years:** Kinrix, Proquad, Lead Hemoglobin + Hematocrit, TB Screening
- **5 years:** No immunizations. Lead, Hemoglobin + Hematocrit
- **6-8 years:** No immunizations
- **9 years:** HPV #1
- **10 years:** HPV #2
- **11 years:** Tdap, Meningococcal ACWY #1
- **12-15 years:** No immunizations
- **16 years:** Meningococcal ACWY #2, Meningococcal B #1*
- **17 years:** Meningococcal B #2*
- **18 years:** No immunizations

Well-Child Care Visits include and examination, review of growth and development information to keep your child healthy, sensory screenings and behavioral assessments, age-appropriate immunizations and lab tests, and discussion of any concerns and questions you may have. Please schedule your child's next visit(s) today at Check-out.



Vaccines missed should be given as soon as possible according to the CDC catch-up schedule.

Influenza vaccine: The CDC recommends everyone 6 months and older get vaccinated every flu season. Children 6 months through 8 years of age receive 2 doses of the influenza vaccine during a single flu season. Everyone else needs only 1 dose each flu season.

***Meningococcal B vaccine:** Bexsero - Administer 2 doses of Bexsero at least 1 month apart. Patients with private insurance only to receive the Meningococcal B vaccine at the pharmacy. There are two Meningococcal B vaccines available. The same vaccine product must be used for all doses.

Rotarix: Administer Rotarix 2 dose series with 1st dose at 2 months old and 2nd dose at 4 months old. The maximum age for the 1st dose is 14 weeks, 6 days old. The maximum age for the 2nd dose is before the patient turns 6 months old.

Vaxelis: Contains DTaP, Hepatitis B, Polio and Hib

Kinrix: Contains DTaP and Polio

ProQuad: Contains Measles, Mumps, Rubella and Varicella

RSV Vaccines for Infants: Neonates and infants born during or entering their first RSV season and children up to 24 months of age who remain vulnerable to severe RSV in their second RSV season should receive RSV vaccine.

Nirsevimab (Beyfortus): Recommended for all infants younger than 8 months of age born during RSV season or entering their first RSV season. Except in rare circumstances, most infants younger than 8 months of age do not need Nirsevimab if they were born 14 or more days after their mother received the RSV vaccine. It is also recommended for some children aged 8-19 months, who are at increased risk for severe RSV disease and entering their second RSV season.

*HPV Schedule:

- Routine vaccination for all adolescents 9 years old through age 18. The number of doses is dependent on age at initial vaccination:
- Age 9-14 years at initiation: 2 dose series at 0 and 6-12 months. Minimum interval: 5 months
- Age 15 years or older at initiation: 3-dose series at 0, 1-2 months, 6 months. Minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months
- Interrupted schedules: If the HPV vaccination schedule is interrupted, the series does not need to be restarted.

Note: CHLAMYDIA screening: 16-24 years.

