

HAVE YOU SCHEDULED YOUR CHILD'S NEXT WELL-CHILD CARE APPOINTMENT?

Visits include examination, review of growth and development, sensory screenings, behavioral assessments, immunizations, lab tests, and discussion of any concerns and questions. Please schedule your child's next visit(s) today at Check-out.

WELL-CHILD VISITS AND IMMUNIZATION SCHEDULE

- **Birth:** Hepatitis B
- **• 3-5 days (for breast-fed newborns):** No immunizations
- **• 1-2 weeks:** No immunizations
- **• 1 month:** No immunizations
- **• 2 months:** DTaP + Polio + Hib + Hepatitis B vaccines #1, PCV-20 #1, Rotavirus #1
- **• 4 months:** DTaP + Polio + Hib + Hepatitis B vaccines #2, PCV-20 #2, Rotavirus #2
- **• 6 months:** DTaP + Polio + Hib + Hepatitis B vaccines #3, PCV-20 #3, Influenza #1
- **• 9 months:** Influenza #2 (if applicable)
- **• 12 months (1 year old):** MMR #1, Varicella #1, Hepatitis A #1
- **• 15 months:** DTaP #4, Hib #4, PCV-20 #4
- **• 18 months:** Hepatitis A #2
- **• 24 months (2 years old):** No immunizations
- **• 30 months (2.5 years old):** No immunizations
- **• 3 years:** No immunizations
- **• 4 years:** DTaP #5, Polio #4, MMR #2, Varicella #2
- **• 5 years:** No immunizations
- **• 6-8 years:** No immunizations
- **• 9 years:** HPV #1
- **• 10 years:** HPV #2
- **• 11 years:** Tdap, Meningococcal ACWY #1
- **• 12-15 years:** No immunizations
- **• 16 years:** Meningococcal ACWY #2, Meningococcal B #1
- **• 17 years:** Meningococcal B #2
- **• 18 years:** No immunizations

See reverse
for more info
on screenings
and vaccines



Additional Immunization Notes

- Everyone **6 months and older** should get an updated **COVID-19 and Flu Vaccine annually**.
- Nirsevimab (RSV Vaccine) is recommended for **babies under 8 months born during or entering their first RSV season. Most do not need if their mother received the vaccine at least 14 days before birth.** It's also recommended for some high-risk children 8–19 months entering their second RSV season. Ask your health care provider if your child qualifies.
- **Patients with private insurance can only receive the Meningococcal B vaccine at the pharmacy.** The same vaccine product must be used for all doses.



SCREENINGS BY AGE

- **Lead, Hemoglobin + Hematocrit:** 9 months, 2-5 years
- **Tuberculosis:** 3-4 years
- **Hearing:** 4-6 years
- **Vision:** 3-6 years
- **Chlamydia:** 16-24 years